

参考文献

<総 説>

1. 夏秋知英 (2009) 弱毒ウイルスの防除機構. (「微生物と植物の相互作用—病害と生物防除—」百町・對馬編) pp134-141. (ISBN978-4-88171-120-0)

<原 著>

2. Miyoshi, H., Okade, H., Muto, S., Suehiro, N., Nakashima, H., Tomoo, K. and Natsuaki, T. (2008). *Turnip mosaic virus VPg interacts with Arabidopsis thaliana eIF(iso)4E and inhibits in vitro translation.* **Biochimie** 90(10): 1427-1434.
3. Goto, K., Kobori, T., Kosaka, Y., Natsuaki, T. and Masuta, C. (2007). Characterization of Silencing Suppressor 2b of Cucumber Mosaic Virus Based on Examination of its Small RNA-Binding Abilities. **Plant and Cell Physiology** 48(7): 1050-1060.
4. Wang, W-Q., Natsuaki, T., Kosaka, Y. and Okuda, S. (2006). Comparison of the nucleotide and amino acid sequences of parental and attenuated isolates of *Zucchini yellow mosaic virus*. **Journal of General Plant Pathology** 72(1): 52-56. (corresponding author)
5. Nakazono-Nagaoka, E., Suzuki, M., Kosaka, Y. and Natsuaki, T. (2005). RT-PCR-RFLP analysis for evaluating cross protection by an attenuated isolate of *Cucumber mosaic virus*. **Journal of General Plant Pathology** 71(3): 243-246.
6. Kobori, T., Ryang, B.-S., Natsuaki, T., and Kosaka, Y. (2005). A new technique to select mild strains of *Cucumber mosaic virus*. **Plant Disease** 89:879-882.
7. Suehiro, N., Natsuaki, T., Watanabe, T. and Okuda, S. (2004). An important determinant of the ability of *Turnip mosaic virus* to infect *Brassica* spp. and/or *Raphanus sativus* is in its P3 protein. **Journal of General Virology** 85(7): 2087-2098. (corresponding author)
8. Nakazono-Nagaoka, E., Sato, C., Kosaka, Y. and Natsuaki, T. (2004). Evaluation for cross-protection of an attenuated isolate of Bean yellow mosaic virus by RT-PCR. **Journal of General Plant Pathology** 70(6):359-362.
9. Kuroda,T., Natsuaki,T., Wang,W.Q. and Okuda,S. (1997). Formation of multimers of cucumber mosaic virus satellite RNA. **Journal of General Virology** 78:941-946. (corresponding author)
10. Sayama,H., Sato, T., Kominato,M., Natsuaki,T. and Kaper,J.M. (1993). Field testing of satellite-containing attenuated strain of cucumber mosaic virus for tomato protection in Japan. **Phytopathology** 83(4):405-410.